

Supplementary Information

Bio-detoxification of ricin in castor bean (*Ricinus communis L.*) seeds

Natália L. Sousa^{1,2}, Glaucia B. Cabral¹, Pabline M. Vieira³, Aisy B. Baldoni^{1,4}, Francisco J. L. Aragão^{1*}


¹ Embrapa Recursos Genéticos e Biotecnologia, PqEB W5 Norte, 70770-900, Brasília, DF, Brazil.

² Universidade de Brasília, Departamento de Biologia Celular, Campus Universitário, 70910-900, Brasília, DF, Brazil.


³ Instituto Federal Goiano, Campus Urutaí, Laboratório de Biotecnologia, 75790-000, Urutaí, GO, Brazil.

⁴ Current address: Embrapa Agrossilvipastoril, Rod. dos Pioneiros MT-222, 78550-970, Sinop, MT, Brazil.

*To whom correspondence may be addressed. Tel: +55 61 34484777; E-mail addresses: francisco.aragao@embrapa.br, fjaragao@gmail.com



Supplementary Figure 1. Small intestine epithelial cells (IEC-6) were exposed to proteins (0 to 50 µg total protein/mL) isolated from endosperm of the transgenic event TB14S-5D and wild type (control). Protein from control plants contains ricin at the concentrations of 0, 1, 5, 10, 25, 50, 100 and 1000 ng ricin/mL. White bar represents 50 µm and magnification is the same in all images.



Supplementary Figure 2. Full-length gel and blot images used to compose the Figure 1.